

Toczeń rumieniowy indukowany lekami

Kod Orpha: 231111 Kod OMIM:

Opis choroby *

Definicja

A rare, systemic disease with skin involvement characterized by the onset of idiopathic lupus erythematosus-like signs and symptoms resulting from continuous drug intake (>1 month), which resolve when treatment is discontinued, in persons with no history of autoimmune disease. Manifestations are variable and may be systemic (e.g. arthralgia, myalgia, fever, fatigue, serositis, pleuritis, pericarditis), subacute cutaneous (incl. photosensitive, non-scarring, annular, polycyclic or papulosquamous lesions, malar erythema, vasculitis, bullous lesions, erythema multiforme-like changes), and/or chronic cutaneous (typically discoid lesions in sun-exposed areas). Procainamide and hydralazine are the drugs most frequently implicated.

Dane

Klasifikacja	Synonimy	
Choroba	DILE	
	DILE	
Kod ORPHA	Kod OMIM	Kod ICD10
231111	-	M32.0
Kod ICD11		
4A40.1		

[*Źródło](#)

orphanet

Rozszerzony opis choroby

Brak opisu rozszerzonego dla tej choroby. Opracowanie w toku.

Orphanet - interntowa baza danych dotyczących rzadkich chorób i sierochych leków. ©INSERM 1999 -
Dostępna na stronie www.orphanet.pl