

# Rak neuroendokrynnny przełyku

Kod Orpha: 506136 Kod OMIM:

## Opis choroby \*

### Definicja

A group of esophageal epithelial neoplasms characterized by neuroendocrine differentiation, comprising well-differentiated neuroendocrine tumors (NETs), poorly differentiated neuroendocrine carcinomas (NECs), and mixed neuroendocrine-non-neuroendocrine neoplasms, an umbrella category including mixed adenoneuroendocrine carcinoma. The tumors typically occur in the lower esophagus, often in association with Barrett mucosa. NECs may also arise in other parts of the esophagus. On endoscopy, NETs usually appear as small polypoid or nodular submucosal masses, while NECs are large, infiltrative, and ulcerated. Patients most commonly present with dysphagia, pain, weight loss, and sometimes melena. Metastatic NETs may be associated with carcinoid syndrome.

### Dane

#### Klasyfikacja

Choroba

#### Synonimy

Esophageal NEN  
Esophageal neuroendocrine neoplasm  
NEN of esophagus  
Esophageal NEN  
Esophageal neuroendocrine neoplasm  
NEN of esophagus

#### Kod ORPHA

506136

#### Kod OMIM

-

#### Kod ICD10

C15.8

#### Kod ICD11

-

---

[\\*Źródło](#)

orphanet

## **Rozszerzony opis choroby**

Brak opisu rozszerzonego dla tej choroby. Opracowanie w toku.

**Orphanet** - interntowa baza danych dotyczących rzadkich chorób i sierochych leków. ©INSERM 1999 -  
Dostępna na stronie [www.orphanet.pl](http://www.orphanet.pl)