

# Nabyty obrzęk naczynioruchowy z niedoborem C1Inh

## Kod Orpha: 528663 Kod OMIM:

### Opis choroby \*

#### Definicja

A rare non-histaminic angioedema characterized by potentially life-threatening episodes of edema of subcutaneous and/or mucosal tissues without urticaria, caused by excessive consumption of C1 esterase inhibitor (C1-INH) in the context of lymphoproliferative or autoimmune diseases. Patients typically present in the fourth decade of life or later and without a family history of angioedema. Clinical manifestation includes nonpitting edema of the skin predominantly involving the face, but also the limbs or genitals, as well as abdominal pain due to involvement of the gastrointestinal mucosa, and severe edema of the upper airway and oral mucosa. Laboratory examination shows low C1-INH activity and low C3, C4, and C1q levels. Autoantibodies to C1-INH are frequently detectable.

#### Dane

#### Klasyfikacja

Choroba

#### Synonimy

Acquired angioneurotic edema with C1 inhibitor deficiency

Nabyty obrzęk angioneurotyczny z niedoborem C1Inh

Nabyty obrzęk angioneurotyczny z niedoborem inhibitora C1

Acquired angioneurotic edema with C1Inh deficiency

#### Kod ORPHA

528663

#### Kod OMIM

-

#### Kod ICD10

D84.1

#### Kod ICD11

-

---

[\\*Źródło](#)

orphanet

## **Rozszerzony opis choroby**

Brak opisu rozszerzonego dla tej choroby. Opracowanie w toku.

**Orphanet** - internetowa baza danych dotyczących rzadkich chorób i sierochych leków. ©INSERM 1999 -  
Dostępna na stronie [www.orphanet.pl](http://www.orphanet.pl)