

## Opis choroby \*

### Definicja

A rare common cystic lymphatic malformation characterized by a benign cystic lesion composed of dilated lymphatic channels. Microcystic lesions consist of cysts smaller than 1 cm in diameter. They usually present at birth or during the first years of life and most often occur in the head and neck region but may affect any site. Symptoms depend on the location and extent of the lesion. Infection, trauma, or intracystic hemorrhage can lead to lesional expansion. Malignant transformation does not occur.

### Dane

#### Klasyfikacja

Zespół wad wrodzonych

#### Synonimy

Capillary lymphangioma  
Malformacja limfatyczna kapilarna  
Malformacja limfatyczna powierzchowna  
Naczyniak limfatyczny kapilarny  
Naczyniak limfatyczny mikrotorbielowaty  
Naczyniak limfatyczny powierzchowny  
Capillary lymphatic malformation  
Cutaneous lymphangioma circumscriptum  
Microcystic infiltrating lymphatic malformation  
Microcystic lymphangioma  
Superficial lymphangioma  
Superficial lymphatic malformation

#### Kod ORPHA

79490

#### Kod OMIM

-

#### Kod ICD10

D18.1

#### Kod ICD11

LA90.11

---

#### [\\*Źródło](#)

orphanet