

Hiperlipidemia z powodu niedoboru wątrobowej lipazy triglicerydowej

Kod Orpha: 140905 Kod OMIM: 614025

Opis choroby *

Definicja

Hyperlipidemia due to hepatic triacylglycerol lipase deficiency is a rare, genetic hyperalphalipoproteinemia disorder characterized by elevated plasma cholesterol and triglyceride (TG) levels with a marked TG enrichment of low- and high-density lipoproteins (HDL), presence of circulating beta-very low density lipoproteins and elevated HDL cholesterol levels, in the presence of a very low, or undetectable, postheparin plasma hepatic lipase activity. Premature atherosclerosis and/or coronary heart disease may be associated.

Dane

Klasyfikacja	Synonimy
Choroba	Hyperlipidemia due to HL deficiency
	Hyperlipidemia due to HTGL deficiency
	Hyperlipidemia due to hepatic lipase deficiency
	Hyperlipidemia due to hepatic triglyceride lipase deficiency

Kod ORPHA	Kod OMIM	Kod ICD10
140905	614025	E78.4

Kod ICD11

-

*[Źródło](#)

orphanet

Rozszerzony opis choroby

Brak opisu rozszerzonego dla tej choroby. Opracowanie w toku.

Orphanet - internetowa baza danych dotyczących rzadkich chorób i sierochych leków. ©INSERM 1999 -
Dostępna na stronie www.orphanet.pl