

# Embriofetopatia antagonistów witaminy K

Kod Orpha: 1914 Kod OMIM:

## Opis choroby \*

### Definicja

Vitamin K antagonist embryofetopathy is characterized by a group of symptoms that may be observed in a fetus or newborn when the mother has taken oral vitamin K antagonists, such as warfarin during pregnancy. Vitamin K antagonists are anticoagulant drugs that provide efficient thromboprophylaxis and that can cross the placenta. 5-12 % of infants exposed to warfarin between 6-9 weeks gestation present nasal hypoplasia and skeletal abnormalities, including short limbs and digits (brachydactyly), and stippled epiphyses. Warfarin fetopathy with central nervous system abnormalities (hydrocephalus, intellectual disability, spasticity, and hypotonia) or ocular abnormalities (microphthalmia, cataract, optic atrophy), fetal loss, and stillbirth, occurs in infants exposed at later gestations. Additional features that have been reported after in utero warfarin exposure include facial dysmorphism (cleft lip and/or palate, malformed ears), choanal atresia or stenosis, aorta coarctation, situs inversus totalis, bilobed lungs, and ventral midline dysplasia.

### Dane

#### Klasyfikacja

Zespół wad wrodzonych

#### Synonimy

Vitamin K antagonist embryopathy  
Zespół di Sala  
Embriopatia warfarynowa  
Embriofetopatia warfarynowa  
Embriopatia antagonistów witaminy K  
Warfarin embryofetopathy  
Warfarin embryopathy  
di Sala syndrome

#### Kod ORPHA

1914

#### Kod OMIM

-

#### Kod ICD10

Q86.2

#### Kod ICD11

LD2F.02

---

[\\*Źródło](#)

orphanet

## **Rozszerzony opis choroby**

Brak opisu rozszerzonego dla tej choroby. Opracowanie w toku.

**Orphanet** - internetowa baza danych dotyczących rzadkich chorób i sierochych leków. ©INSERM 1999 -  
Dostępna na stronie [www.orphanet.pl](http://www.orphanet.pl)